



GE HealthCare

Navigate your journey with clarity and confidence

Welcome to amyloid PET imaging for Alzheimer's assessment, a step toward answers.



VIZAMYL™
Flutemetamol F18
Injection

See Important Safety Information and complete Prescribing Information on the last page.




How Vizamyl supports early detection of Alzheimer's disease

Get the clarity you need to plan for next steps. Rule out Alzheimer's disease or confirm the presence of amyloid plaque to enable clinical decisions, timely interventions, and treatment.

Positron emission tomography (PET) imaging with Vizamyl provides clarity by answering the question of whether amyloid has started to build up in the brain.¹



Vizamyl PET provides pictures that can show where and how much amyloid is in the brain.^{2,3}



With Vizamyl, full-color brain scans show if amyloid has built up in the brain.

How to prepare for amyloid PET imaging with Vizamyl

Before the exam

Tell your doctor if you or the patient who will be scanned:

- are pregnant, may be pregnant or plan to become pregnant
- are breastfeeding
- have any allergies
- have ever had a reaction to Vizamyl or polysorbate 80

If you have any questions about Vizamyl PET imaging, share these with your doctor as well.



How to prepare for amyloid PET imaging with Vizamyl

On the day of the exam



Drink plenty of fluids and use the bathroom frequently.



Wear loose, comfortable clothing.



Ask a family member or caregiver to join you.

What to expect during the exam



The test will be performed by a technologist, a healthcare professional. He or she will inject Vizamyl through a vein in your arm.



A picture of your brain will be taken 1-2 hours later.



To ensure a clear image, lie on the PET camera table as still as possible for 10-20 minutes.


After the exam



Continue drinking plenty of fluids and using the bathroom frequently.



The images of your brain and a report will be sent to your doctor. He or she will explain the test results and what they mean for your next steps.





Support and resources are available for patients and caregivers from Vizamyl. Click here to learn more.



To find imaging centers in your area that are using Vizamyl, click here to access our Vizamyl locator tool.

For more information about Vizamyl, call GE HealthCare Customer Service at 800.292.8514.

To report SUSPECTED ADVERSE REACTIONS, contact GE HealthCare at 800.654.0118 (option 2, then option 1) or by email at GPV.drugsaftey@ge.com or the FDA at 800.FDA.1088 or www.fda.gov/medwatch.



References:

1. Alzforum. FDA approves a second amyloid imaging agent. Available at: www.alzforum.org/news/research-news/fda-approves-second-amyloid-imaging-agent# Accessed on: 04.10.25.
2. Kolanko MA, Win Z, Loreto F, et al. Amyloid PET imaging in clinical practice. *Pract Neurol*. 2020;20(6):451-462.
3. Bollack A, Pemberton HG, Collij LE, et al. Longitudinal amyloid and tau PET imaging in Alzheimer's disease: A systematic review of methodologies and factors affecting quantification. *Alzheimers Dement*. 2023;19(11):5232-5252.

Vizamyl — Important Safety Information for Consumers

Product Use

VIZAMYL™ (flutemetamol F 18 injection) is indicated for positron-emission tomography (PET) of the brain to estimate amyloid beta neuritic plaque density in adults with cognitive impairment for:

- Evaluation of Alzheimer's disease (AD) and other causes of cognitive decline.
- Selection of patients who are indicated for amyloid beta-directed therapy as described in the prescribing information of the therapeutic products.

Important Safety Information About Vizamyl (flutemetamol F 18 injection)

You should not be given Vizamyl if: You have a history of hypersensitivity reaction to Vizamyl or polysorbate 80. Your healthcare professional (HCP) can help you understand what the drug ingredients are.

What is the most important information you should know about Vizamyl?

Allergies: Tell your HCP if you are allergic to any medications or a substance called polysorbate 80.

Reactions such as flushing and shortness of breath have been seen within minutes following administration of Vizamyl. Postmarketing experience included anaphylactic reactions.

Radiation Risk: Vizamyl contributes to your long-term cumulative radiation exposure. Long-term cumulative radiation exposure is associated with an increased risk of cancer. Please be well hydrated and urinate regularly after your test.

The most common side effects of Vizamyl include:

Flushing, increased blood pressure, headache, nausea and dizziness. Instruct patients to inform their healthcare provider if they:

- have had prior reactions to Vizamyl or any component, including polysorbate 80, or
- have reduced renal or hepatic function

Instruct patients to increase their level of hydration before and after receiving Vizamyl (flutemetamol F 18 injection) and to void frequently following Vizamyl administration.

Make sure you tell your HCP:

- All the medications you are taking, including any over-the-counter medications.
- If you are pregnant, may be pregnant or plan to become pregnant.
- If you are breastfeeding.

Children and Elderly:

- Vizamyl should not be used in those less than 18 years old.
- Studies showed no difference in safety between those over 65 years old and those between 18 and 65 years old.

The safety information included here is not comprehensive. If you have any questions, please be sure to discuss them with your HCP.

[Click here for complete Prescribing Information](#)